

**CONTENTS:**

Chlorhexidine Diacetate            2.0%  
Inert Ingredients                        98.0%

**INDICATIONS:**

For use in preparation of a 0.5% chlorhexidine diacetate solution for dipping the teats of cows after milking.

**DIRECTIONS FOR USE:**

To prepare 1 gallon of a 0.5% solution for dipping teats: mix 32 fluid ounces (1 qt.) of Chlorhexidine Diacetate Solution 2% with 6 fluid ounces of glycerin, then add clean potable water until you have a total volume of 1 gallon. Daily teat dipping should start 1 week before cow freshens. During lactation, dip each teat into 0.5% solution immediately after the cow is milked. When the cow is dried off, the teats should continue to be dipped in the 0.5% solution once a day for 3 to 4 days. Dip each teat so that the lower 1 inch of the teat has been covered with the 0.5% solution.



NDC: 50480-122-40



**CHLORHEXIDINE  
DIACETATE**

**2%  
Solution**

TAKE TIME  READ LABEL  
INSTRUCTIONS

**FOR ANIMAL USE ONLY  
KEEP AWAY FROM CHILDREN**

**NET CONTENTS**

1 GALLON (3.79 LITERS)

**STORAGE:**

Store at controlled room temperature between 59°F to 86°F (15°C to 30°C)

**PRECAUTIONS:**

Do not use the concentrate as a teat dip. Hypersensitivity to this product may occur. Use with caution. Udder and teats of the cow should be thoroughly washed before milking. Precipitate may occur with certain types of water used for dilution. It is preferable to use distilled or deionized water. Avoid contamination of animal feed or human ingestion. If contacted, thoroughly flush area with water.

**MANUFACTURED BY:**

Radix Laboratories, Inc.  
1334 International Drive  
Eau Claire, WI 54701  
1-800-927-2349

SEALED FOR YOUR PROTECTION

3.5

6.75